

Year 1 Health Screening Record

Midwifery Education Program MSc in Child Life & Pediatric Psychosocial Care MSc Occupational Therapy Program MSc Physiotherapy Program Nursing Graduate and Undergraduate Programs Physician Assistant Education Program Undergraduate Medicine Program

Congratulations on admission to the Faculty of Health Sciences (FHS) at McMaster University. Completion of a Health Screening Record is a mandatory requirement for registration in your program, and for participation in clinical activities. It can take several weeks to complete so start early! **Review the requirements carefully to ensure accuracy and avoid delay in clinical clearance.**

- The requirements in this Record are in accordance with the Ontario Hospital Association (OHA) Communicable Diseases Screening Protocols and the Council of Ontario Faculties of Medicine (COFM) Immunization Policy. All sections are mandatory except for the suggested requirements on page 5. Exemptions will only be allowed for medical reasons, in which case a note from a physician must be included.
- This Record must be completed by a licensed Health Care Provider (HCP) physician, registered nurse, nurse practitioner. Every HCP who completes any part of this Record must complete the HCP information section on page 2 and initial the applicable sections. HCP initials verify they have either provided the service or they have seen the student's record.
- ** Copies of required laboratory and x-ray reports (if applicable) must be attached. **
- Attach copies of other supporting documentation if available, such as official vaccination records, records from other institutions, or a letter from a physician if applicable HCP signatures/initials for the corresponding sections on this Record are not required. Translate documents into English, if applicable. Make sure your name is on every page. Submit the <u>entire</u> McMaster Record along with your documentation.
- Be sure to complete and sign the student information section on page 2
- If completion of specific requirements is still in process by the deadline for submission, submit completed documentation <u>on time</u> with a note of explanation.
- Once the completed Health Screening Record is reviewed and cleared, a Certificate of Clinical Clearance will be issued.

N.B. Keep the <u>original</u> of this Record and supporting documentation for your files. Documents submitted to the FHS Health Screening Office are not returned.

If you have any questions, please contact the FHS Health Screening Office: 905-525-9140 ext 22249, hrsadmin@mcmaster.ca

Please submit	Please submit by the deadline date below to allow for processing and clearance to:					
FHS HEALTH S	FHS HEALTH SCREENING OFFICE (not your Program Office)					
Preferred:	Submit electronically via secure ShareFile on the McMaster Health Screening website http://fhs.mcmaster.ca/healthscreening/firstyearstudents.html Confidential Fax: 905-528-4348 In Person / Mail <u>copies only</u> : 1280 Main Street West, MDCL 3514, Hamilton ON L8S 4K1 (Emailing personal health information is not recommended)					
Other options:						
Deadline Date	July 15 prior to Year 1	Deadline Date July 31 prior to Year 1				
Physician Assis	te Program se Based s Based	 MSc in Child Life & Pediatric Psychosocial Care MSc Occupational Therapy Program MSc Physiotherapy Program Nursing Undergraduate Program > BScN Accelerated Stream F > BScN Basic Stream A 				

McMaster University values your privacy. Personal information provided on this Record and supporting documentation is protected and is being collected pursuant to the Freedom of Information and Protection of Privacy Act of Ontario (RSO 1990). This information will be held in strict confidence within the Faculty of Health Sciences Health Screening Office and only disclosed as needed with the consent of the student.



STUDENT INFORMATION:		
Name (last):	Name (first):	
Program:	Date of birth:	
Student #: Email:		
(This em	ail address will be used for communication fro	m the FHS Health Screening Office)
I verify that this Record and all supporting document the information provided is accurate.	ation are true copies of the original and t	hat to the best of my knowledge
I understand that it is my responsibility to retain the the duration of my program at McMaster.	priginal of all documentation related to th	e Health Screening Record for
I understand that it is my ethical and professional ob Tuberculosis, Hepatitis B, Hepatitis C and/or HIV.	ligation to inform the Assistant Dean of r	ny Program of any infection with
I understand that failure to complete the requirer in delays and/or removal from participation in cli		new requirements, may result
Signature:	Date:	
HEALTH CARE PROVIDER (HCP) INFORMATION: Every HCP who completes any part of this Record must cc or they have seen the student's record. (Attach additional sh	mplete this section. HCP initials verify the	ney have either provided the service
HCP #1		
Name:	Profession:	Initials:
Address:		
Telephone:	Fax:	
Signature:	Date:	
HCP #2		
Name:	Profession:	Initials:
Address:		
Telephone:	Fax:	
Signature:	Date:	
HCP #3		
Name:	Profession:	Initials:
Address:		
Telephone:	Fax:	
Signature:	Date:	

	in Tests:								
Docum	nent record o	of previous two	-step TB sł	kin test given	at any time ir	the past	(two tests	7-28 days apar	t)
OR If	no record of	previous two-	step TB ski	n test and:		•			
	o tests in last	-	=	o-step TB skir	test require	d (two tes	ts 7-28 da	ys apart)	
	<u>ne</u> test in last							ep One and Ste	ер Тwo
<u>T</u> \	vo tests in las	st 12 months	> Doo	cument both to	ests as Step	One and S	Step Two		
		nust be given						,	
		nust be space		• •	•				
	-	skin tests if tr on is not a cor				ITIVE I B SI	kin test or	active TB disea	se.
					C	mm Indu	ration	Interpretation	HCP Initia
Chan		Date Given do	1/mm/yyyy	Date Read	aa/mm/yyyy	mm Indu		Interpretation	
Step (
Step				. et		I			
Additi	onal TB Skin	test required	after Janua	ary 1 st this yea	r if not includ	led above			
B. Comp	lete if positi [,]	ve TB skin te	st or histo	rv of active T	B disease:				
<u></u>									
		Date Given do	l/mm/yyyy	Date Read	dd/mm/yyyy	mm Indu	ration H	CP Initials	
Positi	ve Test								
Chest	X-Ray requi	ired (must be	subsequer	t to positive te	est) Repo	ort attache	d:		Y
	nt must verify		-		<i>,</i> .			result by a phy	
	<u>Int must venn</u>							to my Progran	
								unexplained we	
					- .				
	:ASLES, M	UMPS, RUE	۶ELLA & ۱	/ΔΚΙ(ΞΕΓΓΔ		ae vaccin		aboratory pro	
tion 2. Mi					: Two dose				oof of immuni
	1970 or later								
• If born			ation (2 dos	ses) is strongl	y recommen	ded over s	serologic t	esting for immu	nity.
If bornSerolog	gic testing for		ation (2 dos	ses) is strongl ended before	y recommen or after MMF	ded over s R vaccinat	serologic to ion. If ser	esting for immu ology is inadve	nity.
If bornSerolog	gic testing for Juent to two N	r immunity is n MMR vaccines	ation (2 dos not recomm s and does	ses) is strongl ended before not demonstra	y recommen or after MMF ate immunity,	ded over s R vaccinat	serologic to ion. If ser	esting for immu ology is inadver ot necessary.	nity.
If bornSerolog	gic testing for Juent to two N Tv	r immunity is n MMR vaccines wo doses vac	ation (2 dos not recomm s and does scine at lea	ses) is strongl ended before not demonstra a st 4 weeks a	y recommen or after MMF ate immunity, part	ded over s R vaccinat , re-vaccir	serologic to ion. If ser nation is no	esting for immu ology is inadve	nity. rtently done
 If born Serolog subsect 	gic testing for Juent to two N Tv	r immunity is n MMR vaccines	ation (2 dos not recomm s and does scine at lea	ses) is strongl ended before not demonstra	y recommen or after MMF ate immunity,	ded over s R vaccinat , re-vaccir	serologic to ion. If ser nation is no Labor	esting for immu ology is inadver ot necessary. <u>OR</u> ratory proof of	nity. rtently done immunity
 If born Serolog subsect 	gic testing for Juent to two N Tv	r immunity is n MMR vaccines wo doses vac	ation (2 dos not recomm s and does scine at lea	ses) is strongl ended before not demonstra a st 4 weeks a	y recommen or after MMF ate immunity, part	ded over s R vaccinat , re-vaccir	serologic to ion. If ser nation is no Labor Measles	esting for immu ology is inadvel of necessary. <u>OR</u> ratory proof of IgG Ab Re	nity. rtently done immunity port attached:
 If born Serolog subsect 	gic testing for Juent to two N Tv	r immunity is n MMR vaccines wo doses vac	ation (2 dos not recomm s and does scine at lea	ses) is strongl ended before not demonstra a st 4 weeks a	y recommen or after MMF ate immunity, part	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u>	serologic to ion. If ser nation is no Labor Measles Mumps lo	esting for immu ology is inadver of necessary. OR ratory proof of IgG Ab Re gG Ab Re	nity. rtently done immunity port attached: port attached:
 If born Serolog subsect 	gic testing for Juent to two N Tv	r immunity is n MMR vaccines wo doses vac	ation (2 dos not recomm s and does scine at lea	ses) is strongl ended before not demonstra a st 4 weeks a	y recommen or after MMF ate immunity, part	ded over s R vaccinat , re-vaccir	serologic to ion. If ser nation is no Labor Measles	esting for immu ology is inadver of necessary. OR ratory proof of IgG Ab Re gG Ab Re	nity. rtently done immunity port attached:
 If born Serolog subsect Measles Mumps Rubella If histo 	gic testing for juent to two M Tv #1 dd/n	r immunity is n MMR vaccines wo doses vac mm/yyyy	ation (2 dos not recomm s and does ccine at lea #2 dd/	ses) is strongl ended before not demonstra ist 4 weeks a /mm/yyyy	y recommend or after MMF ate immunity, part HCP Initials	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u> <u>OR</u>	serologic to ion. If ser hation is no Labor Measles Mumps lo Rubella I uired vario	esting for immu ology is inadver of necessary. <u>OR</u> ratory proof of IgG Ab Re gG Ab Re gG Ab Re	nity. rtently done immunity port attached: port attached: port attached:
 If born Serolog subsect Measles Mumps Rubella If histo Serolog 	gic testing for juent to two M Tv #1 dd/n ry of chicken gic testing for	r immunity is n MMR vaccines wo doses vac mm/yyyy pox or shingle r immunity afte	ation (2 dos not recomm s and does ccine at lea #2 dd/ #2 dd/ es, laborato er Varicella	ses) is strongl ended before not demonstra i st 4 weeks a (mm/yyyy	y recommend or after MMF ate immunity, part HCP Initials HCP Initials	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u> <u>OR</u> urally acq nd not rec	serologic to ion. If ser hation is no Labor Measles Mumps la Rubella I uired vario commende	esting for immu ology is inadvel of necessary. <u>OR</u> ratory proof of IgG Ab Re gG Ab Re gG Ab Re sella/zoster request.	nity. rtently done immunity port attached: port attached: port attached: uired.
 If born Serolog subsect Measles Mumps Rubella If histo Serolog 	gic testing for juent to two M Tv #1 dd/n ry of chicken gic testing for	r immunity is n MMR vaccines wo doses vac mm/yyyy pox or shingle r immunity afte	ation (2 dos not recomm s and does ccine at lea #2 dd/ #2 dd/ es, laborato er Varicella	ses) is strongl ended before not demonstra i st 4 weeks a (mm/yyyy	y recommend or after MMF ate immunity, part HCP Initials HCP Initials	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u> <u>OR</u> urally acq nd not rec	serologic to ion. If ser hation is no Labor Measles Mumps la Rubella I uired vario commende	esting for immu ology is inadver of necessary. <u>OR</u> ratory proof of IgG Ab Re gG Ab Re gG Ab Re	nity. rtently done immunity port attached: port attached: port attached: uired.
 If born Serolog subsect Measles Mumps Rubella If histo Serolog 	gic testing for Juent to two M Tv #1 dd/n gic testing for and Varicella	r immunity is n MMR vaccines wo doses vac mm/yyyy pox or shingle r immunity afte	ation (2 dos not recomm s and does ccine at lea #2 dd/ #2 dd/ es, laborato er Varicella be given a	ses) is strongl ended before not demonstra ist 4 weeks a (mm/yyyy) iry proof of im vaccination is t the same tin	y recommend or after MMF ate immunity, part HCP Initials HCP Initials munity to nat s unreliable a he, otherwise	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u> <u>OR</u> urally acq nd not rec	serologic to ion. If ser hation is no Labor Measles Mumps ly Rubella I uired vario commende to be space	esting for immu ology is inadvel of necessary. OR ratory proof of IgG Ab Re gG Ab Re gG Ab Re gG Ab Re ad at least 4 we <u>OR</u>	nity. rtently done immunity port attached: port attached: port attached: uired. eks apart.
 If born Serolog subsect Measles Mumps Rubella If histo Serolog 	gic testing for Juent to two M Tv #1 dd/n ry of chicken gic testing for and Varicella Tv	r immunity is n MMR vaccines wo doses vac mm/yyyy pox or shingle r immunity afte vaccines may	ation (2 dos not recomm s and does ccine at lea #2 dd/ #2 dd/ es, laborato er Varicella be given a ccine at lea	ses) is strongl ended before not demonstra ist 4 weeks a (mm/yyyy) iry proof of im vaccination is t the same tin	y recommend or after MMF ate immunity, part HCP Initials HCP Initials munity to nat s unreliable a he, otherwise	ded over s R vaccinat , re-vaccir <u>OR</u> <u>OR</u> OR urally acq nd not rec they mus	serologic to ion. If ser hation is no Labor Measles Mumps ly Rubella I uired vario commende to be space	esting for immu ology is inadvel of necessary. OR ratory proof of IgG Ab Re gG Ab Re gG Ab Re gG Ab Re sella/zoster requ ed at least 4 we	nity. rtently done immunity port attached: port attached: port attached: uired. uired.

Section 3. HEPATITIS E	3 (HBV): Primary vaccination	series PLUS Anti-HBs se	erology				
Hepatitis B Primary Vaccination Series (2 dose schedule only if given age 11-15 years)							
	#1 dd/mm/yyyy	#2 dd/mm/yyyy	+/- #3 dd/mm/yyyy	HCP Initials			
HBV primary series	S						
Plus Anti-HBs serolog	Plus Anti-HBs serology (≥ one month > primary series) Report attached: STOP here if ≥ 10 IU/L (immune)						
✤ Anti-HBs after doct	umented primary series < 10 IU/L	(not immune) > One boos	ster dose vaccine required				
	#1 dd/mm/yyyy	HCP Initials					
HBV booster #1							
Repeat Anti-HBs s	Repeat Anti-HBs serology one month > booster #1 Report attached: STOP here if ≥ 10 IU/L (immune)						
✤ Anti-HBs after boos	ster #1 < 10 IU/L (not immune)	Continue two additional b	ooster doses vaccine				
	#2 dd/mm/yyyy	#3 dd/mm/yyyy 5 months > Booster #2	HCP Initials				
HBV boosters #2 8							
Repeat Anti-HBs s	erology one month > booster #3	Report attached:	STOP here if ≥ 10 IU/L (i	mmune)			
 Anti-HBs after boost 	ster #3 < 10 IU/L (not immune) >	> HBs Antigen serology regi	uired Report attached:				
HBs Antigen negat		une. Report status to the Ass	·	n.			
HBs Antigen positiv	•	positive result to the Assista					
Section 4. PERTUSSIS:	Tdan vaccine (Adacel React						
	Tuap vaccille (Adacei, Boost	rix, Repevax, DTCoq)					
	· · ·		dult (≥ age 18 years) health	care workers,			
If currently age 18 years regardless of age, recei	s or older: The Ontario Hospital A ve a single dose of pertussis vaco	ssociation requires that all ac ine (Tdap), if not previously r	eceived in adulthood, ever	n if not due for			
If currently age 18 years regardless of age, recei a tetanus diphtheria boo	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc <i>oster</i> . The interval between the la	ssociation requires that all ac sine (Tdap), if not previously r st tetanus diphtheria booster	eceived in adulthood, ever	n if not due for			
If currently age 18 years regardless of age, recei <i>a tetanus diphtheria boo</i> matter. The adult dose	s or older: The Ontario Hospital A ve a single dose of pertussis vaco	ssociation requires that all ac sine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster.	eceived in adulthood, ever and the adult Tdap vaccine	n if not due for			
If currently age 18 years regardless of age, recei <i>a tetanus diphtheria boo</i> matter. The adult dose If currently < age 18 yea	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1	eceived <u>in adulthood</u> , <i>ever</i> and the adult Tdap vaccine 4-17 years	n if not due for			
If currently age 18 years regardless of age, recei <i>a tetanus diphtheria boo</i> matter. The adult dose If currently < age 18 yea	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles	ssociation requires that all ac sine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster.	eceived in adulthood, ever and the adult Tdap vaccine	n if not due for does not			
If currently age 18 years regardless of age, recei <i>a tetanus diphtheria boo</i> matter. The adult dose If currently < age 18 yea	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1	eceived <u>in adulthood</u> , <i>ever</i> and the adult Tdap vaccine 4-17 years	n if not due for does not			
If currently age 18 years regardless of age, recei <i>a tetanus diphtheria boo</i> matter. The adult dose If currently < age 18 yea Tda	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy	eceived <u>in adulthood</u> , <i>ever</i> and the adult Tdap vaccine 4-17 years <u>Age (years)</u>	n if not due for does not			
If currently age 18 years regardless of age, recei a tetanus diphtheria boo matter. The adult dose If currently < age 18 yea Tda	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years <u>Age (years)</u> oosters if required	HCP Initials			
If currently age 18 years regardless of age, recei a tetanus diphtheria boo matter. The adult dose If currently < age 18 yea Tda Section 5. TETANUS, D • Document record of	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc <i>oster</i> . The interval between the la is <u>in addition</u> to the routine adoles <u>ars</u> : Document record of adolesce up vaccine brand name	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years <u>Age (years)</u> oosters if required Full primary series is 3 vac	HCP Initials			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D • Document record colspan="2">If unable to locate Adult vaccination s	s or older: The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac e childhood vaccination records chedule: Vaccine #2 ≥ 2 months	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – s, you must start a new prin after vaccine #1, Vaccine #3	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) osters if required Full primary series is 3 vac hary series (3 doses) ≥ 6 months after Vaccine #	HCP Initials			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D Occument record colspan="2">If unable to locate Adult vaccination s Require at least on	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac childhood vaccination records ichedule: Vaccine $\#2 \ge 2$ months we polio vaccine age 4 years or old	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – s, you must start a new prin after vaccine #1, Vaccine #3	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) osters if required Full primary series is 3 vac hary series (3 doses) ≥ 6 months after Vaccine #	HCP Initials			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D • Document record colspan="2">If unable to locate Adult vaccination s	<u>s or older</u> : The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac childhood vaccination records ichedule: Vaccine $\#2 \ge 2$ months the polio vaccine age 4 years or old ooster.	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – 5, you must start a new prin after vaccine #1, Vaccine #3 der plus at least one tetanus of	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) oosters if required Full primary series is 3 vacc hary series (3 doses) ≥ 6 months after Vaccine # diphtheria vaccination in las	HCP Initials HCP Initials cines each.			
If currently age 18 years regardless of age, recei a tetanus diphtheria boo matter. The adult dose If currently < age 18 yea Tda Section 5. TETANUS, D • Document record of • If unable to locate Adult vaccination s • Require at least on primary series or b	s or older: The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac e childhood vaccination records ichedule: Vaccine #2 ≥ 2 months the polio vaccine age 4 years or old ooster. #1 dd/mm/yyyy	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – s, you must start a new prin after vaccine #1, Vaccine #3	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) osters if required Full primary series is 3 vac hary series (3 doses) ≥ 6 months after Vaccine #	h if not due for does not HCP Initials			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D Occument record colspan="2">If unable to locate Adult vaccination s Require at least on	s or older: The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac e childhood vaccination records ichedule: Vaccine #2 ≥ 2 months the polio vaccine age 4 years or old ooster. #1 dd/mm/yyyy	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – 5, you must start a new prin after vaccine #1, Vaccine #3 der plus at least one tetanus of	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) oosters if required Full primary series is 3 vacc hary series (3 doses) ≥ 6 months after Vaccine # diphtheria vaccination in las	HCP Initials HCP Initials cines each.			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D Occument record colspan="2">Adult vaccination s Require at least on primary series or b Tetanus, Diphtheri	s or older: The Ontario Hospital A ve a single dose of pertussis vacc oster. The interval between the la is <u>in addition</u> to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima of tetanus, diphtheria and polio vac e childhood vaccination records ichedule: Vaccine #2 ≥ 2 months the polio vaccine age 4 years or old ooster. #1 dd/mm/yyyy	association requires that all ac cine (Tdap), if not previously r st tetanus diphtheria booster scent pertussis booster. ent Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + bo ccinations received to date – 5, you must start a new prin after vaccine #1, Vaccine #3 der plus at least one tetanus of	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) oosters if required Full primary series is 3 vacc hary series (3 doses) ≥ 6 months after Vaccine # diphtheria vaccination in las	HCP Initials HCP Initials cines each.			
If currently age 18 years regardless of age, receir a tetanus diphtheria boo matter. The adult dose If currently < age 18 years Tda Tda Section 5. TETANUS, D Occument record colspan="2">Adult vaccination s Require at least on primary series or b Tetanus, Diphtheri	s or older: The Ontario Hospital A ve a single dose of pertussis vaccoster. The interval between the la is in addition to the routine adoles is in addition to the routine adoles ars: Document record of adolesce up vaccine brand name IPHTHERIA & POLIO: Prima Prima of tetanus, diphtheria and polio vace childhood vaccination records ichedule: Vaccine #2 ≥ 2 months up polio vaccine age 4 years or old ooster. #1 dd/mm/yyyy a +/- #4 dd/mm/yyyy a	Association requires that all action (Tdap), if not previously rest tetanus diphtheria booster scent pertussis booster. Tdap vaccine given age 1 Date dd/mm/yyyy ary vaccination series + booster ccinations received to date – 5, you must start a new prin after vaccine #1, Vaccine #3 der plus at least one tetanus of #2 dd/mm/yyyy	eceived <u>in adulthood</u> , ever and the adult Tdap vaccine 4-17 years Age (years) oosters if required Full primary series is 3 vacc hary series (3 doses) ≥ 6 months after Vaccine # diphtheria vaccination in las #3 dd/mm/yyyy	HCP Initials HCP Initials Lines each. HCP Initials HCP Initials			

SUGGESTED REQUIREMENTS:
The following are <u>not</u> requirements of the Faculty of Health Sciences at McMaster University; <u>however</u> one or more may be mandatory for some elective or clinical placements.
□ Influenza – Vaccination with current season's vaccine by December 1 st each year strongly recommended
Meningitis – Men-C-ACWY vaccination (Menactra)
Hepatitis A vaccination – Recommended for travel to endemic areas
□ Polio – One booster dose vaccine ≥ age 18 years recommended for travel to countries where poliomyelitis is prevalent
 Blood Borne Viruses – <u>strongly recommended</u> Hepatitis B (HBV) – HBV Surface Antigen (HBsAg) serology Hepatitis C (HCV) – HCV Antibody serology HIV – HIV Antibody serology N.P. Studente who are infected with Hepatitie P. Hepatitie C and/or HIV/ must cell report their statue to the Assistant Deep
N.B. Students who are infected with Hepatitis B, Hepatitis C and/or HIV must self-report their status to the Assistant Dean of their program.